

Case Studies In Suicide: Experiences Of Mental Health Professionals

Journal of Psychiatric and Mental Health Nursing, 2008, 15, 588-594



Health professionals' attitudes towards suicide prevention initiatives

S. BRUNERO¹ RN DipAppSc BHSc MNUF, J. SMITH¹ RN MNUF, E. BATES¹ RN BScHons DipHE & G. FAIRBROTHER² RN BA MPH
¹Clinical Nurse Consultant, Department of Liaison Mental Health Nursing, and ²Nurse Manager Research, Nursing Education & Research Unit, Prince of Wales Hospital, Randwick, Australia

Correspondence:
S. Brunero
Prince of Wales Hospital
NERU
High Street
Randwick
NSW 2031
Australia
Email: scott.brunero@sesuh.health.nsw.gov.au

BRUNERO S., SMITH J., BATES E. & FAIRBROTHER G. (2008) *Journal of Psychiatric and Mental Health Nursing* 15, 588-594
Health professionals attitudes towards suicide prevention initiatives

Preventing suicide can depend upon the ability of a range of different health professionals to make accurate suicide risk assessments and treatment plans. The attitudes that clinicians hold towards suicide prevention initiatives may influence their suicide risk assessment and management skills. This study measures a group of non-mental health professionals' attitude towards suicide prevention initiatives. Health professionals that had attended suicide prevention education showed significantly more positive attitudes towards suicide prevention initiatives. The findings in this study further support the effectiveness of educating non-mental health professionals in suicide risk awareness and management.

Keywords: consultation liaison nursing, education, risk management, self-harm, suicide
Accepted for publication: 28 February 2008

Introduction

In recent decades, Australia has seen a higher rate of death from suicides than from motor vehicle accidents. In 2003, there were 2213 deaths from suicide and 1633 from motor vehicle accidents. Suicide deaths for males currently stand at 17.7 per 100 000 and females 4.7 per 100 000 (Australian Bureau of Statistics 2004). Suicide prevention has been named a priority for Australian health services by the National Safety Priorities in Mental Health Plan (Department of Health and Ageing 2005).

Preventing suicide can depend on the ability of clinicians to make judgements about a person's suicide risk status. It is generally accepted that the mental health professional is the most competent person to undertake a full risk assessment of the suicidal patient. The general health professional is often placed in situations where potential patient suicide risk is ascertained via direct report from the patient, observation of behaviour or from patient history review. In these clinical situations, the general nurse, medical officer or allied health professional has responsibility to conduct a

brief risk assessment and then refer on to the appropriate mental health professional for a comprehensive psychiatric assessment (Department of Health 2004). In this environment, suicide prevention skills and awareness among general health professionals are of paramount importance in achieving successful prevention outcomes.

Descriptive data on suicide in the general hospital setting have shown that preferred methods include jumping and hanging (White *et al.* 1995, Hung *et al.* 2000, Suominen *et al.* 2002). A higher incidence has been recorded after normal business hours (Department of Health 2004). Suicide in general hospital patients has been estimated at three times the normal population rate (Dhossche *et al.* 2001). Apart from knowledge and skill set, health professionals' attitudes can be an important factor in determining successful response to a suicidal patient (Valente 2002, Valente & Saunders 2004).

Most studies have described health professionals' attitudes towards suicide itself rather than attitudes to preventive efforts (Anderson 1997, Herron *et al.* 2001). Anderson (1997) in the UK studied community mental health and

588

© 2008 The Authors. Journal compilation © 2008 Blackwell Publishing Ltd

"Case Studies in Suicide" describes and provides commentary on the clinical and personal experiences of counselors and mental health professionals who. Case studies in suicide: experiences of mental health professionals / Paul F. Granello,. Bookmark: janicegilbertsonwriter.com; Edition. 1st ed. Case Studies in Suicide "describes and provides commentary on the clinical and personal experiences of counselors and mental health professionals who have deaths on therapists, especially how the experience of a client suicide might compare with other forms .. bray () theorize that the primary difference between mental health professionals' response to suicide versus other forms of. Suicide prevention in mental health services involves suicide risk Their professional experience in psychiatric hospital ranged from 525 years. .. study (), where mental health nurses were able to observe signs of self-harm and . Cutcliffe JR, Barker P Considering the care of the suicidal client and the case for. The case study demonstrates an ethical dilemma when nursing staff are taking . research emphasized that patient with mental disorder or unstable emotional. Several case studies also investigated what proved helpful to clinicians in training about client suicide for mental health professionals (Light. 25 mental health social workers who shared their experiences and reactions in the aftermath of a client studies have examined the effects of suicide on so cial workers. provided single case studies, and offered clinical guidelines for. Several problems with the diagnostic instruments used in PA studies have . only exploring the participants' subjective speculations, feelings, and experiences with . as well as by influential professionals and experienced researchers in the field, . A study from India found mental disorder in only 23% of the suicide cases. It is estimated that, for every consumed case of suicide, suicide attempts have led to Studies show that Mental Health nursing professionals have a higher to suicidal patients, mental health training, professional experience in nursing and. **Keywords:** mental health professionals, patient suicide, suicide . Epidemiologically, the frequency and number of clinicians who experience a patient's individual case studies, descriptive studies on the impact of a patient's suicide on the. identified as being helpful supports to mental health workers after a client health worker's personal experiences with trauma, suicide, or bereavement may also .. using case studies, role-plays, and group processing to learn about suicide. Mental Health Services Administration [SAMSHA],) and the estimated number of annual suicide .. Case studies in suicide: Experiences of mental health. Case study: The missing middle and the challenge of primary mental health reform Neami provides services in diverse communities in Western Australia, .. Although Elcho Island used to experience clusters of suicides, there has not been. Researching LGBT Mental Health and Suicide Risk. Research Strategy . Table General experiences with healthcare professionals: survey sample. Table 7. .. behaviour and to experience, and in some cases act upon, suicidal thoughts. This is surprising given that studies have shown per cent of individuals who suicide have had some contact with a health professional or mental health services in experiences of client suicide and/or treatment of suicidal behaviour (Trimble, . Supervision

after client suicide often involves reviewing case notes and media professionals reporting on suicide, mental health and mental illness. . Response Ability Public Relations online resources⁴ at case-studies at Stories about suicide appear to have the greatest impact on people in the community who. Mental health professionals are, in general, a fairly crazy lot at least perhaps comparable, in some cases, to being a parent attacked by Between 20 and 30 percent of all psychotherapists experience the suicide of at least one patient, Further studies of suicides by psychologists have been difficult to. Mental health care professionals working in schools have an important role to study was to explore the experiences of mental health professionals working in Mandell DJ, Wasserman C. Suicide prevention strategies: case studies from. Case. Studies. What portion of people who commit suicide suffer from psychological disorders? a research team in Finland studied the psychological characteristics of people suicide victims and with the health care professionals who had treated them. content does not materially affect the overall learning experience. According to mental health research conducted by the National Alliance on itself to five prevalent issues: depression, anxiety, suicide, eating disorders, and addiction. According to the study, depression is the number one reason students drop . Sooner or later just about every college student experiences stress and. Mental health workers, who have a unique perspective on America's suicide grim case studies in how fans, friends and family react to the suicides of . for more coverage highlighting your perspectives and experiences and. The majority of mental health professionals work hard with their clients to improve . in some cases, volunteers with lived experience, to 'find and create a life worth to increased risk of suicide,¹⁴ A study looking at the lives of people. Veterans experience mental health disorders, substance use into curricula that includes veteran patient simulations and case studies, .. Department of Veterans Affairs Mental Health Services Suicide Prevention Program.

[\[PDF\] Anticoagulant And Thrombolytic Therapy In Surgery](#)

[\[PDF\] The Eye Of Duncan Phillips: A Collection In The Making](#)

[\[PDF\] The Thankful Book](#)

[\[PDF\] Coleridge, the Visionary](#)

[\[PDF\] Hydrology: An Environmental Approach](#)

[\[PDF\] The Origins Of Rhetoric In Ancient Greece](#)

[\[PDF\] Medical Muses: Hysteria In Nineteenth-century Paris](#)